

SCHEDULING STATUS

WARNING: (A) PREMATURE DISCONTINUATION OF XABANAR INCREASES THE RISK OF THROMBOTIC EVENTS, (B) SPINAL/EPIDURAL HAEMATOMA

Premature discontinuation of XABANAR increase the risk of thrombotic events:
Premature discontinuation of any oral anticoagulant, including XABANAR, increases the risk of thrombotic events. If anticoagulation with XABANAR is discontinued for a reason other than the pathological bleeding or completion of a course of therapy, consider coverage with another anticoagulant [see Posology and method of administration (4.2), and Special Warnings and precautions for use (4.4)].

Spinal/epidural haematoma:
Epidural or spinal haematoma have occurred in patients treated with XABANAR who are receiving neuraxial anaesthesia or undergoing spinal puncture. These haematomas may result in feeling numb or paralysed or in permanent paralysis.

Consider these risks when scheduling the patients for spinal procedures.

Factors that increase the risk of developing epidural or spinal haematomas in these patients include:

- Use of indwelling epidural catheters.
- Concomitant use of other medicines that affect haemostasis, such as nonsteroidal anti-inflammatory drugs (NSAIDs), platelet inhibitors, other anticoagulants.
- History of spinal deformity or spinal surgery.
- Concomitant use of neuraxial anaesthesia and/or XABANAR and neuraxial procedures is not known to increase the risk of spinal haematoma (4.4) and **undetectable effects (4.8)**.

Monitor patients carefully for signs and symptoms of neurological impairment. If neurological impairment occurs, urgent treatment is necessary [see **Special Warnings and Precautions (4.4)**]. Consider the benefits and risks before neuraxial intervention in patients anticoagulated or to be anticoagulated for thromboprophylaxis [see **Special Warnings and precautions (4.4)**].

1. NAME OF THE MEDICINE

XABANAR 10 mg film-coated tablets

XABANAR 15 mg film-coated tablets

XABANAR 20 mg film-coated tablets

XABANAR 10 mg tablets contain 10 mg of rivaroxaban (creatinine clearance < 50 > 150 ml/min). Contains sugar lactose monohydrate 26.0 mg per tablet.

XABANAR 15 mg tablets contain 15 mg of rivaroxaban (creatinine clearance < 30 – 15 ml/min). Contains sodium 0.59 mg per tablet.

XABANAR 20 mg tablets contain 20 mg of rivaroxaban (creatinine clearance < 30 – 15 ml/min). Contains sodium 0.53 mg per tablet.

XABANAR 20 mg: Each film-coated tablet contains rivaroxaban 20 mg.

Contains sugar lactose monohydrate 19.50 mg per tablet.

Contains sodium 0.53 mg per tablet.

For full list of excipients, see section 6.1

2. PHARMACOTHERAPEUTIC FORM

Film-coated tablet.

XABANAR 10 mg: Round, light red, biconvex film-coated tablets having "A1" debossed on one side and plain on other side.

XABANAR 15 mg: Round, red, biconvex film-coated tablets having "A2" debossed on one side and plain on other side.

XABANAR 20 mg: Dark red coloured, oval shaped, biconvex film coated tablet with "A3" debossed on one side and plain on other side.

4.1. CLINICAL PARTICULARS

4.1. Therapeutic indications: XABANAR is indicated for the prevention of venous thromboembolism (VTE) in patients undergoing major orthopaedic surgery of the lower limbs.

XABANAR 15 mg and XABANAR 20 mg are indicated for the prevention of non-valvular atrial fibrillation (SPAF).

• Treatment of deep vein thrombosis (DVT) and for the prevention of recurrent deep vein thrombosis (DVT) and pulmonary embolism (PE).

• Treatment of postoperative embolism (PE) and for the prevention of recurrent pulmonary embolism (PE) and postoperative thrombosis (DVT).

4.2. Posology and method of administration

Recommended dose and frequency of administration

The recommended dose is taken with XABANAR 10 mg tablet once daily for the prevention of venous thromboembolism (VTE) in patients undergoing major orthopaedic surgery.

The initial dose should be taken within 6 - 10 hours after surgery provided that haemostasis has been established.

If a dose is missed during the 15 mg twice daily treatment phase (day 1 to 21), the patient should take XABANAR immediately to ensure intake of 30 mg XABANAR per day. In this case two 15 mg tablets may be taken at once. The patient should continue with the regular 15 mg twice daily intake as recommended after the missed day.

If a dose is missed during the once daily treatment phase, the patient should take XABANAR immediately, and continue on the following day with the once daily intake as recommended. The dose should not be doubled within the same day to make up for a missed dose.

SPAF - Maximum daily dose

The recommended maximum daily dose is on XABANAR 20 mg tablet (20 mg rivaroxaban).

SPAF - Additional information on special populations

SPAF - Patients with hepatic impairment

XABANAR 15 mg and XABANAR 20 mg are contraindicated in patients with hepatic disease with Child Pugh B and C. Limited clinical data are available for patients with severe hepatic impairment (Child Pugh B and C) (see section 4.3 and 5.2).

Limited clinical data in patients with moderate hepatic impairment (Child Pugh B) indicate a significant increase in the pharmacological activity.

Use of XABANAR 15 mg are available for patients with severe hepatic impairment (Child Pugh C) (see sections 4.3 and 5.2).

SPAF - Patients with renal impairment

No dose adjustment is required if XABANAR 10 mg is administered in patients with mild (creatinine clearance < 30 to 50 ml/min) renal impairment. For patients with moderate (creatinine clearance < 30 to 15 ml/min) renal impairment the recommended dose is one XABANAR 15 mg once daily.

Limited clinical data for patients with severe renal impairment (creatinine clearance < 30 to 15 ml/min) indicate that rivaroxaban plasma levels are significantly increased in this patient population. Therefore XABANAR 10 mg must be used with caution in these patients (see section 4.4). Use is not recommended in patients with severe renal impairment (creatinine clearance < 15 ml/min) (see sections 4.4 and 5.2).

Ethnic differences

No dose adjustment is required based on ethnic factors (see section 4.5).

XABANAR 15 mg and XABANAR 20 mg

There is no need for monitoring of coagulation parameters during treatment with XABANAR 15 mg and XABANAR 20 mg.

SPAF - Additional information on special populations

The recommended dose is on XABANAR 20 mg tablet once daily.

For patients with moderate renal impairment (creatinine clearance < 50 to 30 ml/min) the recommended dose is one XABANAR 15 mg tablet once daily.

For patients with severe renal impairment (creatinine clearance < 30 ml/min) the recommended dose is one XABANAR 10 mg tablet once daily.

Therapy should be continued as long as risk factors for stroke and systemic embolism persist.

SPAF - Missed dose

If a dose is missed the patient should take XABANAR 15 mg or XABANAR 20 mg immediately and continue with the once daily intake as recommended on the following day. The dose should not be doubled to make up for a missed dose within the same day.

SPAF - Maximum daily dose

The recommended maximum daily dose is on XABANAR 20 mg tablet (20 mg rivaroxaban).

SPAF - Additional information on special populations

SPAF - Patients with hepatic impairment

XABANAR 15 mg and XABANAR 20 mg are contraindicated in patients with hepatic disease with Child Pugh B and C. Limited clinical data are available for patients with severe hepatic impairment (Child Pugh B and C) (see section 4.3 and 5.2).

Limited clinical data in patients with moderate hepatic impairment (Child Pugh B) indicate a significant increase in the pharmacological activity.

Use of XABANAR 15 mg are available for patients with severe hepatic impairment (Child Pugh C) (see sections 4.3 and 5.2).

SPAF - Patients with renal impairment

No dose adjustment is required if XABANAR 10 mg is administered in patients with mild (creatinine clearance < 30 to 50 ml/min) renal impairment. For patients with moderate (creatinine clearance < 30 to 15 ml/min) renal impairment the recommended dose is one XABANAR 15 mg once daily.

Limited clinical data for patients with severe renal impairment (creatinine clearance < 30 to 15 ml/min) indicate that rivaroxaban plasma levels are significantly increased in this patient population. Therefore XABANAR 10 mg must be used with caution in these patients (see section 4.4). Use is not recommended in patients with severe renal impairment (creatinine clearance < 15 ml/min) (see sections 4.4 and 5.2).

Ethnic differences

No dose adjustment is required based on ethnic factors (see section 4.5).

XABANAR 15 mg and XABANAR 20 mg

There is no need for monitoring of coagulation parameters during treatment with XABANAR 15 mg and XABANAR 20 mg.

SPAF - Additional information on special populations

The recommended dose is on XABANAR 20 mg tablet once daily.

For patients with moderate renal impairment (creatinine clearance < 50 to 30 ml/min) the recommended dose is one XABANAR 15 mg tablet once daily.

For patients with severe renal impairment (creatinine clearance < 30 ml/min) the recommended dose is one XABANAR 10 mg tablet once daily.

Therapy should be continued as long as risk factors for stroke and systemic embolism persist.

SPAF - Missed dose

If a dose is missed the patient should take XABANAR 15 mg or XABANAR 20 mg immediately and continue with the once daily intake as recommended on the following day. The dose should not be doubled to make up for a missed dose within the same day.

SPAF - Maximum daily dose

The recommended maximum daily dose is on XABANAR 20 mg tablet (20 mg rivaroxaban).

SPAF - Additional information on special populations

SPAF - Patients with hepatic impairment

XABANAR 15 mg and XABANAR 20 mg are contraindicated in patients with hepatic disease with Child Pugh B and C. Limited clinical data are available for patients with severe hepatic impairment (Child Pugh B and C) (see section 4.3 and 5.2).

Limited clinical data in patients with moderate hepatic impairment (Child Pugh B) indicate a significant increase in the pharmacological activity.

Use of XABANAR 15 mg are available for patients with severe hepatic impairment (Child Pugh C) (see sections 4.3 and 5.2).

SPAF - Patients with renal impairment

No dose adjustment is required if XABANAR 10 mg is administered in patients with mild (creatinine clearance < 30 to 50 ml/min) renal impairment. For patients with moderate (creatinine clearance < 30 to 15 ml/min) renal impairment the recommended dose is one XABANAR 15 mg once daily.

Limited clinical data for patients with severe renal impairment (creatinine clearance < 30 to 15 ml/min) indicate that rivaroxaban plasma levels are significantly increased in this patient population. Therefore XABANAR 10 mg must be used with caution in these patients (see section 4.4). Use is not recommended in patients with severe renal impairment (creatinine clearance < 15 ml/min) (see sections 4.4 and 5.2).

Ethnic differences

No dose adjustment is required based on ethnic factors (see section 4.5).

XABANAR 15 mg and XABANAR 20 mg

There is no need for monitoring of coagulation parameters during treatment with XABANAR 15 mg and XABANAR 20 mg.

SPAF - Additional information on special populations

The recommended dose is on XABANAR 20 mg tablet once daily.

For patients with moderate renal impairment (creatinine clearance < 50 to 30 ml/min) the recommended dose is one XABANAR 15 mg tablet once daily.

For patients with severe renal impairment (creatinine clearance < 30 ml/min) the recommended dose is one XABANAR 10 mg tablet once daily.

Therapy should be continued as long as risk factors for stroke and systemic embolism persist.

SPAF - Missed dose

If a dose is missed the patient should take XABANAR 15 mg or XABANAR 20 mg immediately and continue with the once daily intake as recommended on the following day. The dose should not be doubled to make up for a missed dose within the same day.

SPAF - Maximum daily dose

The recommended maximum daily dose is on XABANAR 20 mg tablet (20 mg rivaroxaban).

SPAF - Additional information on special populations

SPAF - Patients with hepatic impairment

XABANAR 15 mg and XABANAR 20 mg are contraindicated in patients with hepatic disease with Child Pugh B and C. Limited clinical data are available for patients with severe hepatic impairment (Child Pugh B and C) (see section 4.3 and 5.2).

Limited clinical data in patients with moderate hepatic impairment (Child Pugh B) indicate a significant increase in the pharmacological activity.

Use of XABANAR 15 mg are available for patients with severe hepatic impairment (Child Pugh C) (see sections 4.3 and 5.2).

SPAF - Patients with renal impairment

No dose adjustment is required if XABANAR 10 mg is administered in patients with mild (creatinine clearance < 30 to 50 ml/min) renal impairment. For patients with moderate (creatinine clearance < 30 to 15 ml/min) renal impairment the recommended dose is one XABANAR 15 mg once daily.

Limited clinical data for patients with severe renal impairment (creatinine clearance < 30 to 15 ml/min) indicate that rivaroxaban plasma levels are significantly increased in this patient population. Therefore XABANAR 10 mg must be used with caution in these patients (see section 4.4). Use is not recommended in patients with severe renal impairment (creatinine clearance < 15 ml/min) (see sections 4.4 and 5.2).

Ethnic differences

No dose adjustment is required based on ethnic factors (see section 4.5).

XABANAR 15 mg and XABANAR 20 mg

There is no need for monitoring of coagulation parameters during treatment with XABANAR 15 mg and XABANAR 20 mg.

SPAF - Additional information on special populations

The recommended dose is on XABANAR 20 mg tablet once daily.

For patients with moderate renal impairment (creatinine clearance < 50 to 30 ml/min) the recommended dose is one XABANAR 15 mg tablet once daily.

For patients with severe renal impairment (creatinine clearance < 30 ml/min) the recommended dose is one XABANAR 10 mg tablet once daily.

Therapy should be continued as long as risk factors for stroke and systemic embolism persist.

SPAF - Missed dose

If a dose is missed the patient should take XABANAR 15 mg or XABANAR 20 mg immediately and continue with the once daily intake as recommended on the following day. The dose should not be doubled to make up for a missed dose within the same day.

SPAF - Maximum daily dose

The recommended maximum daily dose is on XABANAR 20 mg tablet (20 mg rivaroxaban).

SPAF - Additional information on special populations

SPAF - Patients with hepatic impairment

XABANAR 15 mg and XABANAR 20 mg are contraindicated in patients with hepatic disease with Child Pugh B and C. Limited clinical data are available for patients with severe hepatic impairment (Child Pugh B and C) (see section 4.3 and 5.2).

Limited clinical data in patients with moderate hepatic impairment (Child Pugh B) indicate a significant increase in the pharmacological activity.

Use of XABANAR 15 mg are available for patients with severe hepatic impairment (Child Pugh C) (see sections 4.3 and 5.2).

SPAF - Patients with renal impairment

No dose adjustment is required if XABANAR 10 mg is administered in patients with mild (creatinine clearance < 30 to 50 ml/min) renal impairment. For patients with moderate (creatinine clearance < 30 to 15 ml/min) renal impairment the recommended dose is one XABANAR 15 mg once daily.

Limited clinical data for patients with severe renal impairment (creatinine clearance < 30 to 15 ml/min) indicate that rivaroxaban plasma levels are significantly increased in this patient population. Therefore XABANAR 10 mg must be used with caution in these patients (see section 4.4). Use is not recommended in patients with severe renal impairment (creatinine clearance < 15 ml/min) (see sections 4.4 and 5.2).

Ethnic differences

No dose adjustment is required based on ethnic factors (see section 4.5).

XABANAR 15 mg and XABANAR 20 mg

There is no need for monitoring of coagulation parameters during treatment with XABANAR 15 mg and XABANAR 20 mg.

SPAF - Additional information on special populations

The recommended dose is on XABANAR 20 mg tablet once daily.

For patients with moderate renal impairment (creatinine clearance < 50 to 30 ml/min) the recommended dose is one XABANAR 15 mg tablet once daily.

For patients with severe renal impairment (creatinine clearance < 30 ml/min) the recommended dose is one XABANAR 10 mg tablet once daily.

Therapy should be continued as long as risk factors for stroke and systemic embolism persist.

SPAF - Missed dose

If a dose is missed the patient should take XABANAR 15 mg or XABANAR 20 mg immediately and continue with the once daily intake as recommended on the following day. The dose should not be doubled to make up for a missed dose within the same day.

SPAF - Maximum daily dose

The recommended maximum daily dose is on XABANAR 20 mg tablet (20 mg rivaroxaban).

SPAF - Additional information on special populations

SPAF - Patients with hepatic impairment

XABANAR 15 mg and XABANAR 20 mg are contraindicated in patients with hepatic disease with Child Pugh B and C. Limited clinical data are available for patients with severe hepatic impairment (Child Pugh B and C) (see section 4.3 and 5.2).

Limited clinical data in patients with moderate hepatic impairment (Child Pugh B) indicate a significant increase in the pharmacological activity.

Use of XABANAR 15 mg are available for patients with severe hepatic impairment (Child Pugh C) (see sections 4.3 and 5.2).

SPAF - Patients with renal impairment

No dose adjustment is required if XABANAR 10 mg is administered in patients with mild (creatinine clearance < 30 to 50 ml/min) renal impairment. For patients with moderate (creatinine clearance < 30 to 15 ml/min) renal impairment the recommended dose is one XABANAR 15 mg once daily.

Limited clinical data for patients with severe renal impairment (creatinine clearance < 30 to 15 ml/min) indicate that rivaroxaban plasma levels are significantly increased in this patient population. Therefore XABANAR 10 mg must be used with caution in these patients (see section 4.4). Use is not recommended in patients with severe renal impairment (creatinine clearance < 15 ml/min) (see sections 4.4 and 5.2).

Ethnic differences

No dose adjustment is required based on ethnic factors (see section 4.5).

XABANAR 15 mg and XABANAR 20 mg

There is no need for monitoring of coagulation parameters during treatment with XABANAR 15 mg and XABANAR 20 mg.

SPAF - Additional information on special populations

The recommended dose is on XABANAR 20 mg tablet once daily.

For patients with moderate renal impairment (creatinine clearance < 50 to 30 ml/min) the recommended dose is one XABANAR 15 mg tablet once daily.

For patients with severe renal impairment (creatinine clearance < 30 ml/min) the recommended dose is one XABANAR 10 mg tablet once daily.

Therapy should be continued as long as risk factors for stroke and systemic embolism persist.

SPAF - Missed dose

If a dose is missed the patient should take XABANAR 15 mg or XABANAR 20 mg immediately and continue with the once daily intake as recommended on the following day. The dose should not be doubled to make up for a missed dose within the same day.

SPAF - Maximum daily dose

The recommended maximum daily dose is on XABANAR 20 mg tablet (20 mg rivaroxaban).

SPAF - Additional information on special populations

SPAF - Patients with hepatic impairment

XABANAR 15 mg and XABANAR 20 mg are contraindicated in patients with hepatic disease with Child Pugh B and C. Limited clinical data are available for patients with severe hepatic impairment (Child Pugh B and C) (see section 4.3 and 5.2).

Limited clinical data in patients with moderate hepatic impairment (Child Pugh B) indicate a significant increase in the pharmacological activity.

Use of XABANAR 15 mg are available for patients with severe hepatic impairment (Child Pugh C) (see sections 4.3 and 5.2).

SPAF - Patients with renal impairment

No dose adjustment is required if XABANAR 10 mg is administered in patients with mild (creatinine clearance < 30 to 50 ml/min) renal impairment. For patients with moderate (creatinine clearance < 30 to 15 ml/min) renal impairment the recommended dose is one XABANAR 15 mg once daily.

Limited clinical data for patients with severe renal impairment (creatinine clearance < 30 to 15 ml/min) indicate that rivaroxaban plasma levels are significantly increased in this patient population. Therefore XABANAR 10 mg must be used with caution in these patients (see section 4.4). Use is not recommended in patients with severe renal impairment (creatinine clearance < 15 ml/min) (see sections 4.4 and 5.2).

Ethnic differences

No dose adjustment is required based on ethnic factors (see section 4.5).

XABANAR 15 mg and XABANAR 20 mg

There is no need for monitoring of coagulation parameters during treatment with XABANAR 15 mg and XABANAR 20 mg.

SPAF - Additional information on special populations

The recommended dose is on XABANAR 20 mg tablet once daily.

For patients with moderate renal impairment (creatinine clearance < 50 to 30 ml/min) the recommended dose is one XABANAR 15 mg tablet once daily.

For patients with severe renal impairment (creatinine clearance < 30 ml/min) the recommended dose is one XABANAR 10 mg tablet once daily.

Therapy should be continued as long as risk factors for stroke and systemic embolism persist.

SPAF - Missed dose

If a dose is missed the patient should take XABANAR 15 mg or XABANAR 20 mg immediately and continue with the once daily intake as recommended on the following day. The dose should not be doubled to make up for a missed dose within the same day.

SPAF - Maximum daily dose

The recommended maximum daily dose is on XABANAR 20 mg tablet (20 mg rivaroxaban).

SPAF - Additional information on special populations

SPAF - Patients with hepatic impairment

XABANAR 15 mg and XABANAR 20 mg are contraindicated in patients with hepatic disease with Child Pugh B and C. Limited clinical data are available for patients with severe hepatic impairment (Child Pugh B and C) (see section 4.3 and 5.2).

Limited clinical data in patients with moderate hepatic impairment (Child Pugh B) indicate a significant increase in the pharmacological activity.

Use of XABANAR 15 mg are available for patients with severe hepatic impairment (Child Pugh C) (see sections 4.3 and 5.2).

SPAF - Patients with renal impairment

No dose adjustment is required if XABANAR 10 mg is administered in patients with mild (creatinine clearance < 30 to 50 ml/min) renal impairment. For patients with moderate (creatinine clearance < 30 to 15 ml/min) renal impairment the recommended dose is one XABANAR 15 mg once daily.